

COMPONENT I	RECOMMENDATIONS
Hand hygiene	After touching blood, body fluids, secretions, excretions, contaminated items; immediately after removing gloves; between patient contacts.
Personal protective equipment (PPE)	
Gloves	For touching blood, body fluids, secretions, excretions, contaminated items; for touching mucous membranes and nonintact skin
Gown	During procedures and patient-care activities when contact of clothing/exposed skin with blood/body fluids, secretions, and excretions is anticipated..
Mask, eye protection (goggles), face shield*	During procedures and patient-care activities likely to generate splashes or sprays of blood, body fluids, secretions, especially suctioning, endotracheal intubation
Soiled patient-care equipment	Handle in a manner that prevents transfer of microorganisms to others and to the environment; wear gloves if visibly contaminated; perform hand hygiene.
Environmental control	Develop procedures for routine care, cleaning, and disinfection of environmental surfaces, especially frequently touched surfaces in patient-care areas.
Textiles and laundry	Handle in a manner that prevents transfer of microorganisms to others and to the environment
Needles and other sharps	Do not recap, bend, break, or hand-manipulate used needles; if recapping is required, use a one-handed scoop technique only; use safety features when available; place used sharps in puncture-resistant container
Patient resuscitation	Use mouthpiece, resuscitation bag, other ventilation devices to prevent contact with mouth and oral secretions

Patient placement	<p>Prioritize for single-patient room if patient is at increased risk of transmission, is likely to contaminate the environment, does not maintain appropriate hygiene, or is at increased risk of acquiring infection or developing adverse outcome following infection.</p>
<p>Respiratory hygiene/cough etiquette (source containment of infectious respiratory secretions in symptomatic patients, beginning at initial point of encounter e.g., triage and reception areas in emergency departments and physician offices)</p>	<p>Instruct symptomatic persons to cover mouth/nose when sneezing/coughing; use tissues and dispose in no-touch receptacle; observe hand hygiene after soiling of hands with respiratory secretions; wear surgical mask if tolerated or maintain spatial separation, >3 feet if possible.</p>

* During aerosol-generating procedures on patients with suspected or proven infections transmitted by respiratory aerosols (e.g., SARS), wear a fit-tested N95 or higher respirator in addition to gloves, gown, and face/eye protection.